

Europ. J. Combinatorics (1996) **17**, 741–749

Complete Sequences in \mathbb{N}^2

NORBERT HEGYVÁRI

Folkman's theorem states that if $A = \{a_1 < a_2 < \dots\}$ satisfying $A(n) > n^{\frac{1}{2}+\varepsilon}$, where $A(n) = \sum_{a_i \leq n} 1$, then $P(A) = \{a_{i_1} + a_{i_2} + \dots + a_{i_s}; i_1 < i_2 < \dots < i_s; s \in \mathbb{N}; a_{i_j} \in A\}$ contains an infinite arithmetic progression. We shall consider the analogue of Folkman's theorem in \mathbb{N}^2 , and some related questions are also investigated.

© 1996 Academic Press Limited

1. INTRODUCTION

In 1966, J. Folkman proved that if $A = \{a_1 < a_2 < \dots < a_n \dots\}$ is a strictly increasing sequence of integers satisfying $A(n) > N^{\frac{1}{2}+\alpha}$ for some $\alpha > 0$ (where $A(n) = \sum_{a_i \leq n} 1$), then the set $P(A) = \{\sum \varepsilon_i a_i \mid \varepsilon_i = 0 \text{ or } 1; \sum \varepsilon_i < \infty\}$ contains an infinite arithmetic progression. Let us call such an A subcomplete. Furthermore, if $P(A)$ intersects every arithmetic progression, then $P(A)$ contains all sufficiently large integers. Let us say then that A is complete.

In this paper we are going to investigate the analogous question in \mathbb{N}^2 .

We shall prove that the analogue theorem of Folkman's theorem is not true, and we give some other results.

2. NOTATIONS AND DEFINITIONS

Let us use the following notation. Let $\mathbf{v} \in \mathbb{N}^2$; $\mathbf{v} = (v_1, v_2)$, $v_1, v_2 \in \mathbb{N}$. We shall sometimes write $v^{(1)}(v^{(2)})$ for $v_1(v_2)$. Let

$$m(\mathbf{v}) = \begin{cases} v_2/v_1 & \text{if } v_1 \neq 0, \\ \infty & \text{if } v_1 = 0. \end{cases}$$

Let $\mathbf{v}_1, \mathbf{v}_2 \in \mathbb{N}^2$ and let

$$\mathbf{v}_1 + \mathbf{v}_2 = (v_1^{(1)} + v_2^{(1)}, v_1^{(2)} + v_2^{(2)})$$

For $\mathbf{A}, \subseteq \mathbb{N}^2$, denote by $\mathbf{A} + \mathbf{B} = \{\mathbf{x} \mid \mathbf{x} = \mathbf{a} + \mathbf{b} \text{ where } \mathbf{a} \in \mathbf{A} \text{ and } \mathbf{b} \in \mathbf{B}\}$. Let $\mathbf{v} \in \mathbb{N}^2$. Denote by $|\mathbf{v}|$ the length of \mathbf{v} , i.e. $|\mathbf{v}| = \sqrt{v_1^2 + v_2^2}$. Let

$$\mathbf{S}_n = \{\mathbf{v} \mid \mathbf{v} \in \mathbb{N}^2 \text{ and } |\mathbf{v}| \leq n\}.$$

Let $\mathbf{A} = \{\mathbf{a}_1, \mathbf{a}_2, \dots\} \subseteq \mathbb{N}^2$ and let $\mathbf{A}^{(n)} = \{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\}$. We define the counting function of \mathbf{A} as $\mathbf{A}(n) = |\mathbf{A} \cap \mathbf{S}_n|$, where $|\mathbf{X}|$ is the cardinality of a finite set \mathbf{X} .

An analogous notion of an interval is the discrete rectangle. Let

$$\mathbf{R}(a, b, c, d) = \{\mathbf{v} \mid \mathbf{v} \in \mathbb{N}^2, a \leq v_1 \leq c \text{ and } b \leq v_2 \leq d\}$$

and let $t(\mathbf{R})$ be the size of $\mathbf{R}(a, b, c, d)$, i.e. $t(\mathbf{R}) = (c - a + 1)(d - b + 1)$. For given $\mathbf{x}, \mathbf{d} \in \mathbb{N}^2$, call the set

$$L(\mathbf{x}_0, \mathbf{d}) = \{\mathbf{x}_0 + k\mathbf{d} \mid k \in \mathbb{N}\}$$

an arithmetic progression. Let $L(\mathbf{d}) := L(\mathbf{0}, \mathbf{d})$. For a given set $\mathbf{A} = \{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n, \dots\} \subseteq \mathbb{N}^2$, let

$$P(\mathbf{A}) = \left\{ \sum \varepsilon_i \mathbf{a}_i \mid \varepsilon_i = 0 \text{ or } 1; \sum \varepsilon_i < \infty \right\}.$$

Let us call \mathbf{A} L -complete if there exist $\mathbf{x}_0, \mathbf{d} \in \mathbb{N}^2$ such that $L(\mathbf{x}_0, \mathbf{d}) \subset P(\mathbf{A})$.

We can define another completeness, as well. Let us call $\mathbf{A} \subseteq \mathbb{N}^2$ p -complete if there exist $\mathbf{x}_0 \in \mathbb{N}^2$, $\mathbf{u}, \mathbf{v} \in \mathbb{R}^2$, $m(\mathbf{u}) \neq m(\mathbf{v})$, such that

$$S(\mathbf{x}_0, \mathbf{u}, \mathbf{v}) := \{\mathbf{x}_0 + \alpha \mathbf{u} + \beta \mathbf{v} \mid \alpha, \beta \in \mathbb{R}^+ \} \cap \mathbb{N}^2 \subset P(\mathbf{A}).$$

3. THE PLANE ANALOGUE OF FOLKMAN'S THEOREM

In [2], the following remarkable result is established:

THEOREM F (Folkman). *If A is an infinite set of positive integers such that there exist $\varepsilon > 0$ and n_0 with*

$$A(n) > n^{\frac{1}{2} + \varepsilon}$$

for $n > n_0$, then A is subcomplete.

In this section we prove that the analogous theorem does not hold for a sequence $\mathbf{A} \subseteq \mathbb{N}^2$.

THEOREM 3.1. *There exist an $\alpha > 0$ and an $\mathbf{A} \subseteq \mathbb{N}^2$ for which*

$$\mathbf{A}(n) > \alpha n^2$$

and \mathbf{A} is not L -complete.

PROOF OF THEOREM 3.1. Let $0 < \gamma < \pi/4$ and $\varepsilon = \tan \gamma$. Let

$$\mathbf{R}_{2n-1} := \{\mathbf{x} \mid \mathbf{x} \in \mathbb{N}^2; 0 < m(\mathbf{x}) < \varepsilon \text{ and } 2^{4^{2n-1}} \leq |\mathbf{x}| < 2^{4^{2n}}\}$$

and

$$\mathbf{R}_{2n} := \{\mathbf{x} \mid \mathbf{x} \in \mathbb{N}^2; 1/\varepsilon < m(\mathbf{x}) < \infty \text{ and } 2^{4^{2n}} \leq |\mathbf{x}| < 2^{4^{2n+1}}\}$$

$n = 1, 2, \dots$

Now let $\mathbf{A} := \bigcup_{n=1}^{\infty} \mathbf{R}_n$.

First we prove that $\mathbf{A}(n) > \alpha n^2$ for some $\alpha > 0$, $n \in \mathbb{N}$. To prove it, let

$$(y_1, y_2) \in \mathbf{R}'_{2n} \text{ iff } (y_2, y_1) \in \mathbf{R}_{2n}, \quad n \in \mathbb{N}.$$

Clearly, $|\mathbf{R}_{2n}| = |\mathbf{R}'_{2n}|$ for every $n \in \mathbb{N}$. Furthermore,

$$\bigcup_{n=1}^{\infty} (\mathbf{R}_{2n-1} \cup \mathbf{R}'_{2n}) = \{\mathbf{x} \mid \mathbf{x} \in \mathbb{N}^2; 0 < m(\mathbf{x}) < \varepsilon\},$$

so we have

$$\mathbf{A}(N) = |\mathbf{A} \cap \mathbf{S}_N| = \left| \bigcup_{n=1}^{\infty} (\mathbf{R}_{2n-1} \cup \mathbf{R}'_{2n}) \cap \mathbf{S}_N \right| \geq (\gamma/2) |\mathbf{S}_N| / (\pi/2) = \alpha N^2.$$

Now we show that $P(\mathbf{A})$ does not contain an arithmetic progression. Let us assume that, contrary to the assertion, there are $\mathbf{x}_0, \mathbf{d} \in \mathbb{N}^2$ such that $L(\mathbf{x}_0, \mathbf{d}) \subset P(\mathbf{A})$. Let $|\mathbf{d}| = d$.

We distinguish two cases: $0 \leq m(\mathbf{d}) < 1$ or $1 \leq m(\mathbf{d}) \leq \infty$.

The proofs for the cases $0 \leq m(\mathbf{d}) \leq 1$ and $1 \leq m(\mathbf{d}) \leq \infty$ are similar. Thus, without loss of generality, let $0 \leq m(\mathbf{d}) \leq 1$ and $0 \leq m(\mathbf{x}_0) \leq 1$.

LEMMA 3.1. *Let*

$$\mathbf{E}_n := \{\mathbf{x} \mid \mathbf{x} \in \mathbb{N}^2; 4 \cdot \sqrt{2} \cdot 2^{3 \cdot 4^{2n}} / (1 - \varepsilon) < |\mathbf{x}| < 2^{4^{2n+1}} \text{ and } 0 \leq m(\mathbf{x}) \leq 1\}, \quad n \in \mathbb{N}.$$

Then $\mathbf{E}_n \not\subset P(\mathbf{A})$.

PROOF OF LEMMA 3.1. Let

$$\mathbf{F}_n := \{\mathbf{x} \mid \mathbf{x} \in \mathbb{N}^2; 2^{4^{2n}} < |\mathbf{x}| < 2^{4^{2n+1}} \text{ and } 0 \leq m(\mathbf{x}) < 1\}$$

and let $\mathbf{x} \in \mathbf{F}_n \cap P(\mathbf{A})$.

If $\mathbf{a}_i \in \bigcup_{k=n}^{\infty} \mathbf{R}_{2k+1}$, then $|\mathbf{a}_i| > 2^{4^{2n+1}}$. Thus, writing

$$\mathbf{x} = \mathbf{a}_{i_1} + \mathbf{a}_{i_2} + \cdots + \mathbf{a}_{i_s},$$

we have $\mathbf{a}_{i_j} \in \mathbf{U} \cup \mathbf{V}$, where $\mathbf{U} := \bigcup_{k=1}^n \mathbf{R}_{2k-1}$ and $\mathbf{V} := \bigcup_{k=1}^{\infty} \mathbf{R}_{2k}$, which implies

$$\mathbf{F}_n \cap P(\mathbf{A}) \subset P(\mathbf{U} \cup \mathbf{V}) = P(\mathbf{U}) + P(\mathbf{V}).$$

If $\mathbf{z} \in P(\mathbf{U})$; $\mathbf{z} = \mathbf{a}_{j_1} + \mathbf{a}_{j_2} + \cdots + \mathbf{a}_{j_t}$ then, by the triangle inequality, we obtain

$$|\mathbf{z}| \leq \sum_{r=1}^t |\mathbf{a}_{j_r}| \leq \left| \bigcup_{k=1}^n \mathbf{R}_{2k-1} \right| \cdot \max |\mathbf{a}_{j_r}| < 4 \cdot 2^{2 \cdot 4^{2n}} \cdot 2^{4^{2n}} = 4 \cdot 2^{3 \cdot 4^{2n}}. \quad (3.1.1)$$

Now let

$$\mathbf{K} := \{\mathbf{x} \mid \mathbf{x} \in \mathbb{N}^2; 1/\varepsilon \leq m(\mathbf{x}) < \infty\}.$$

Clearly, $P(\mathbf{V}) \subset \mathbf{K}$, so by (3.1.1) we have

$$\mathbf{F}_n \cap P(\mathbf{A}) \subset \{\mathbf{x} \mid 0 \leq m(\mathbf{x}) < \varepsilon; |\mathbf{x}| < 4 \cdot 2^{3 \cdot 4^{2n}}\} + \mathbf{K}. \quad (3.1.2)$$

This implies by an easy calculation that $2^{4^{2n+1}} > |\mathbf{x}| > 4 \cdot \sqrt{2} \cdot 2^{3 \cdot 4^{2n}} / (1 - \varepsilon)$ and $0 \leq m(\mathbf{x}) \leq 1$. Then $\mathbf{x} \notin \mathbf{F}_n \cap P(\mathbf{A})$, which proves the lemma. \blacksquare

Since

$$\Delta_n := 2^{4^{2n+1}} - 4 \cdot \sqrt{2} \cdot 2^{3 \cdot 4^{2n}} / (1 - \varepsilon) \rightarrow \infty$$

if $n \rightarrow \infty$, there is an n_0 such that, for $n > n_0$,

$$\Delta_n > d.$$

Let us write $\mathbf{x}_k = \mathbf{x}_0 + k \cdot \mathbf{d} = (x_{k1}, x_{k2})$. Then, for every k , $0 \leq m(\mathbf{x}_k) \leq 1$ and $x_{k+1,1} - x_{k,1} < d$, respectively. We obtain that, for every $n > n_0$, there is a k such that $\mathbf{x}_k \in \mathbf{E}'_n \not\subset P(\mathbf{A})$. This contradiction proves the theorem. \square

4. SEQUENCES WITH BOUNDED GAPS

If A is a sequence of positive integers, then there is an obvious sufficient condition for its completeness: if $A = A_1 \cup A_2$, $A_1 \cap A_2 = \emptyset$, $P(A_1)$ has bounded gaps (i.e. there is a $K > 0$ such that if $P(A_1) = \{p_1 < p_2 < \cdots\}$, then, for every n , $p_{n+1} - p_n < K$), and $P(A_2)$ contains sufficiently large intervals then A is complete. What happens if $A \subset \mathbb{N}$ is replaced by $\mathbf{A} \subset \mathbb{N}^2$?

DEFINITION 4.1. We say that $\mathbf{X} \subseteq \mathbb{N}^2$ has bounded gaps if there is a sequence $\{\mathbf{x}_i\}$ of

distinct elements of \mathbf{X} such that the differences $\mathbf{x}_{i+1} - \mathbf{x}_i$ assume only finitely many different values.

We prove the following theorem.

THEOREM 4.1. *There is an $\mathbf{A} \subseteq \mathbb{N}^2$ for which $\mathbf{A} = \mathbf{A}_1 \cup \mathbf{A}_2$; $\mathbf{A}_1 \cap \mathbf{A}_2 = \emptyset$ and:*

(1) *for every $r, s \in \mathbb{N}$, there are $a, b \in \mathbb{N}$ such that*

$$\mathbf{R}(a, b, a + r, b + s) \subset P(\mathbf{A}_1); \quad (4.1)$$

(2) *$P(\mathbf{A}_2)$ has bounded gaps;
and \mathbf{A} is not L -complete.*

This \mathbf{A} will have the following property:

$$\mathbf{a}_{n+1} - \mathbf{a}_n \in \{(1, 0), (0, 1)\} \quad \text{for every } n \in \mathbb{N}. \quad (4.2)$$

We say that a sequence \mathbf{A} is simple if (4.2) holds.

We mention that simple sequences have many interesting combinatorial properties (see [1] and [3]). Although a simple sequence \mathbf{A} is not always complete, the set $P(\mathbf{A})$ contains a large regular part.

DEFINITION 4.2. Let $\mathbf{A} \subseteq \mathbb{N}^2$ be a simple sequence and let

$$T_A(n) := \max\{t(\mathbf{R}) \mid \mathbf{R} \text{ is a discrete rectangle and } \mathbf{R} \subset P(\mathbf{A}^{(n)})\}.$$

If $\mathbf{a}_1 = (1, 2)$ and $\mathbf{a}_{n+1} - \mathbf{a}_n = (1, 0)$ for $n \geq 1$, then it is not hard to see that $T_A(n) = n$.

Let $\mathbf{A} \subseteq \mathbb{N}^2$ and let us assume that

$$|\{i : \mathbf{a}_{i+1} - \mathbf{a}_i = (1, 0)\}| = |\{j : \mathbf{a}_{j+1} - \mathbf{a}_j = (0, 1)\}| = \infty. \quad (4.3)$$

We shall prove that if (4.3) holds, then $T(n)$ is much larger than n .

THEOREM 4.2. (1) *Let $\mathbf{A} \subseteq \mathbb{N}^2$ be a simple sequence. If \mathbf{A} satisfies (4.3) and n is large enough, then*

$$n^{\frac{3}{2}}/360 < T_A(n).$$

(2) *On the other hand, there is a simple sequence \mathbf{A} which satisfies (4.3) and*

$$T_A(n) < 4n^2.$$

Finally in this section, we give a sufficient condition for the p -completeness of $\mathbf{A} \subseteq \mathbb{N}^2$.

THEOREM 4.3. *Let $\mathbf{X} \subseteq \mathbb{N}^2$. Suppose that there are $\mathbf{d}_1, \mathbf{d}_2 \in \mathbb{N}^2$, $m(\mathbf{d}_1) \neq m(\mathbf{d}_2)$ and $\varepsilon > 0$, $\varepsilon \in \mathbb{R}$, for which*

$$|L(\mathbf{d}_i) \cap \mathbf{X} \cap \mathbf{S}_n| > n^{\frac{1}{2} + \varepsilon}, \quad (4.4)$$

for $i = 1, 2$.

Assume that $\mathbf{A} := \mathbf{X} \setminus (L(\mathbf{d}_1) \cup L(\mathbf{d}_2))$ satisfies (4.3). Then \mathbf{X} is p -complete.

First, we need some preliminary lemmas.

DEFINITION 4.3. We write

$$\mathcal{H}(B, S, A) := \{(u, v) \mid v = A; B < u \leq B + S\},$$

$$\mathcal{V}(A, S, B) := \{(u, v) \mid u = A; B < v \leq B + S\},$$

where $A, B, S \in \mathbb{N}$, and let

$$P_2(\mathbf{B}) := \{x_2 \mid \mathbf{x} \in P(\mathbf{B}), \mathbf{x} = (x_1, x_2), \mathbf{B} \subseteq \mathbb{N}^2\}.$$

Let

$$H(n) := |\{i: 2 \leq i \leq n; \mathbf{a}_i = \mathbf{a}_{i-1} + (1, 0)\}|,$$

$$V(n) := |\{j: 2 \leq j \leq n; \mathbf{a}_j = \mathbf{a}_{j-1} + (0, 1)\}|.$$

LEMMA 4.1. Let \mathbf{A} be an arbitrary sequence of \mathbb{N}^2 . Let $\mathbf{A}' = \{\mathbf{a}_{i_1}, \mathbf{a}_{i_2}, \dots, \mathbf{a}_{i_t}\}$ be a (finite) subsequence of \mathbf{A} . Suppose that for each j , $1 \leq j \leq t$, we have $\mathbf{a}_{i_j} - \mathbf{a}_{i_{j-1}} = (1, 0)$ (or $(0, 1)$). Then there are $A, B \in \mathbb{N}$ such that $P(\mathbf{A}) \supset \mathcal{H}(A, t/2, B)$ (or $P(\mathbf{A}) \supset \mathcal{V}(A, t/2, B)$).

PROOF OF LEMMA 4.1. Let $\mathbf{A}'' := \{\mathbf{a}_{i_{2j}} \mid \mathbf{a}_{i_{2j}} \in \mathbf{A}', j = 1, 2, \dots, \lfloor t/2 \rfloor\}$. Let $\mathbf{L}_1 := \{\mathbf{a}_{i_1}\}$; $A_1 = a_{i_1}^{(2)}$, $B_1 = a_{i_1}^{(1)}$ and $D_1 = 1$ and, for $1 \leq j \leq \lfloor t/2 \rfloor$, let

$$\mathbf{L}_j := \{L_{j-1} + \mathbf{a}_{i_{2j}}\} \cup \{\mathbf{L}_{j-1} + \mathbf{a}_{i_{2j-1}}\}$$

and $A_j = A_{j-1} + a_{i_{2j-1}}^{(2)}$; $B_j = B_{j-1} + a_{i_{2j-1}}^{(1)}$ and $D_j = D_{j-1} + 1$.

We claim that, for each j , $1 \leq j \leq \lfloor t/2 \rfloor$,

$$\mathcal{H}(B_j, D_j, A_j) \subset \mathbf{L}_j. \quad (4.5)$$

For $j = 1$, (4.5) is just the definition. Let $j > 1$. If $(u, v) \in \mathbf{L}_{j-1}$ then, by

$$(\mathbf{a}_{i_{2j}} + (u, v)) - (\mathbf{a}_{i_{2j-1}} + (u, v)) = (1, 0),$$

we obtain

$$\mathbf{L}_j \supset \mathcal{H}(B_{j-1} + a_{i_{2j-1}}^{(1)}, D_{j-1} + 1, A_{j-1} + a_{i_{2j-1}}^{(2)}) = \mathcal{H}(B_j, D_j, A_j).$$

Let $j = \lfloor t/2 \rfloor$. Then $D_{\lfloor t/2 \rfloor} = \lfloor t/2 \rfloor$ and thus

$$P(\mathbf{A}) \supset \mathbf{L}_{\lfloor t/2 \rfloor} \supset \mathcal{H}(B_{\lfloor t/2 \rfloor}, \lfloor t/2 \rfloor, A_{\lfloor t/2 \rfloor}). \quad \square$$

LEMMA 4.2. Let $H_1, H_2 \subset \mathbb{N}^2$; $\mathbf{H}_1 \cap \mathbf{H}_2 = \emptyset$. Suppose that

$$\mathcal{H}(A_1, D_1, B_1) \subset P(\mathbf{H}_1) \text{ and } \mathcal{V}(A_2, D_2, B_2) \subset P(\mathbf{H}_2).$$

Then there are $A, B \in \mathbb{N}$, such that

$$\mathbf{R}(A, B, A + D_1, B + D_2) \subset P(\mathbf{H}_1 \cup \mathbf{H}_2).$$

PROOF OF LEMMA 4.2. Let $A = A_1 + A_2$; $B = B_1 + B_2$. If $0 < u \leq D_1$ and $0 < v \leq D_2$, then $\mathbf{x}_1 = (A_1 + u, B_1) \in \mathcal{H}(A_1, D_1, B_1)$ and $\mathbf{x}_2 = (A_2, B_2 + v) \in \mathcal{V}(A_2, D_2, B_2)$. Thus

$$\mathbf{x}_1 + \mathbf{x}_2 = (A_1 + A_2 + u, B_1 + B_2 + v) = (A + u, B + v) \in P(\mathbf{H}_1) + P(\mathbf{H}_2) = P(\mathbf{H}_1 \cup \mathbf{H}_2).$$

But

$$\mathbf{R}(A, B, A + D_1, B + D_2) = \{\mathbf{x}_1 + \mathbf{x}_2 \mid 0 < u \leq D_1 \text{ and } 0 < v \leq D_2\},$$

which completes the proof. \square

PROOF OF THEOREM 4.1. Let $\mathbf{a}_1 = (0, 0)$ and let us write

$$\mathbf{a}_n = \mathbf{a}_{n-1} + \begin{cases} (0, 1) & \text{if } n = 2^k, \\ (1, 0) & \text{if } n \neq 2^k. \end{cases} \quad (4.6)$$

Let $\mathbf{A}_1 = \{\mathbf{a}_k \mid k \equiv 0 \text{ or } 3 \pmod{4}\}$ and $\mathbf{A}_2 = \mathbf{A} \setminus \mathbf{A}_1$.

Clearly, \mathbf{A}_1 satisfies (4.3); thus by Lemmas 4.1 and 4.2 we obtain (4.1). The sequence \mathbf{A}_2 (and hence $P(\mathbf{A}_2)$) has bounded gaps. Indeed,

$$\mathbf{a}_{4k+1} - \mathbf{a}_{4k-2} = (\mathbf{a}_{4k+1} - \mathbf{a}_{4k}) + (\mathbf{a}_{4k} - \mathbf{a}_{4k-1}) + (\mathbf{a}_{4k-1} - \mathbf{a}_{4k-2}) = \mathbf{c}_1 + \mathbf{c}_2 + \mathbf{c}_3$$

and $\mathbf{c}_i \in \{(0, 1), (1, 0)\}$ for $i = 1, 2, 3$.

We are now going to prove that \mathbf{A} is not L -complete.

Assume that, contrary to the assertion, there are $\mathbf{x}_0, \mathbf{d} \in \mathbb{N}^2$ such that $L(\mathbf{x}_0, \mathbf{d}) \subset P(\mathbf{A})$. Observing that $\lim_{k \rightarrow \infty} a_k^{(2)} = \infty$, we have that, for each $k \in \mathbb{N}$,

$$|\{\mathbf{x} : \mathbf{x} \in P(\mathbf{A}), x^{(2)} < k\}| < \infty.$$

This implies that $m(\mathbf{d}) > 0$.

We claim that for every $\varepsilon > 0$ there is an $n_0 \in \mathbb{N}$ such that if $\mathbf{x} \in P(\mathbf{A})$ and $x^{(1)} > n_0$, then $m(\mathbf{x}) < \varepsilon$. Hence, choosing ε less than $m(\mathbf{d})$, we obtain a contradiction.

By (4.6) we have that $\mathbf{a}_n = (n - \lfloor \log_2 n \rfloor, \lfloor \log_2 n \rfloor)$. Thus there is an $n_1 \in \mathbb{N}$ such that $m(\mathbf{a}_n) < \varepsilon/2$ for every $n > n_1$. Let $n_1 > \max\{2, 2/\varepsilon\}$ and let $n_0 = n_1^3$.

Let $\mathbf{x} \in P(\mathbf{A})$, $x^{(1)} > n_0$, and let us write \mathbf{x} in the form

$$\mathbf{x} = \sum_{a_i \in A} \varepsilon_i \mathbf{a}_i = \sum_1 \varepsilon_i \mathbf{a}_i + \sum_2 \varepsilon_i \mathbf{a}_i$$

($\varepsilon_i = 0$ or 1) where \sum_1 contains those \mathbf{a}_i for which $a_i^{(1)} \leq n_1$ and \sum_2 contains those ones for which $a_i^{(1)} > n_1$. Let $\mathbf{x}_1 = \sum_1 \varepsilon_i \mathbf{a}_i$ and $\mathbf{x}_2 = \sum_2 \varepsilon_i \mathbf{a}_i$. Then, by (4.6), we obtain

$$x_1^{(1)} < 2 \cdot \binom{n_1}{2} < n_1^2. \quad (4.7)$$

Clearly, $m(\mathbf{x}_1) \leq 1$. Furthermore,

$$m\left(\sum_2 \varepsilon_i \mathbf{a}_i\right) < \varepsilon/2. \quad (4.8)$$

Indeed, since $m(\mathbf{a}_i) < \varepsilon/2$, i.e. $a_i^{(2)} < (\varepsilon/2)a_i^{(1)}$, we have $\sum_2 \varepsilon_i a_i^{(2)} < (\varepsilon/2) \sum_2 \varepsilon_i a_i^{(1)}$, which implies that $x_2^{(2)} < (\varepsilon/2)x_2^{(1)}$, i.e. $m(\mathbf{x}_2) < \varepsilon/2$. Now, by (4.7) and (4.8), we have

$$m(\mathbf{x}) = x^{(2)}/x^{(1)} \leq \{(x^{(1)} - n_1^2) \cdot (\varepsilon/2) + n_1^2\}/x^{(1)} < \varepsilon/2 + n_1^2/x^{(1)} < \varepsilon/2 + 2/n_1 < \varepsilon,$$

since $n_1 > 2/\varepsilon$. \square

PROOF OF THEOREM 4.2. Let $k := \lfloor \sqrt{n}/6 \rfloor$. We can assume without loss of generality that $H := H(n) - H(k) > n/3$ and $a_1^{(2)} > 10$. Now we need a lemma:

LEMMA 4.3. Let $\mathbf{A} \subseteq \mathbb{N}^2$ be a simple sequence. Suppose that \mathbf{A} satisfies (4.3). Then there are $B, n_0 \in \mathbb{N}$ such that, for $n > n_0$,

$$[B + 1, B + a_1^{(2)}\sqrt{n}/40] \subset P_2(\mathbf{A}^{(k)}).$$

PROOF OF LEMMA 4.3. By (4.3) we can assume that $V(k) > \max\{10a_1^{(2)}, 5(a_1^{(2)} + 1)/(a_1^{(2)} - 5)\}$. Let $D := \lfloor V(k)/5 \rfloor$ and $z := \min\{P \mid V(P) = D\}$. Let us write $P_2(\mathbf{A}^{(z)}) = \{x_1 < x_2 < \dots < x_s\}$. Clearly, $s > z \geq D$. Since \mathbf{A} is a simple sequence, and by the definition of z , we have $a_z^{(2)} = a_1^{(2)} + D$. This implies that, for every i , $1 \leq i < s$,

$$x_{i+1} - x_i \leq a_1^{(2)} + D. \quad (4.9)$$

Since $V(k) - V(z) \geq 4D$, by Lemma 4.1, we can select a subset $\mathbf{A}^* = \{\mathbf{a}_{i_1}, \mathbf{a}_{i_1-1}, \dots, \mathbf{a}_{i_{2D}}, \mathbf{a}_{i_{2D}-1} \mid \mathbf{a}_{i_j} - \mathbf{a}_{i_{j-1}} = (0, 1); \mathbf{a}_{i_{j-1}} \neq \mathbf{a}_{i_{j+1}} \text{ for } j = 1, \dots, 2D\}$ of $\{\mathbf{a}_{z+1}, \mathbf{a}_{z+2}, \dots, \mathbf{a}_k\}$ such that $P(\mathbf{A}^*) \supset \mathcal{V}(A, 2D, B)$ for some $A, B \in \mathbb{N}$.

In view of (4.9) and by $D > a_1^{(2)}$, we obtain that there is a $T \in \mathbb{N}$ such that

$$P_2(\mathbf{A}^{(z)}) + P_2(\mathbf{A}^*) \supset [T + 1, T + (x_s - x_1)]. \quad (4.10)$$

Since $a_i^{(2)} \geq a_1^{(2)}$, we have

$$x_s - x_1 \geq a_1^{(2)} \cdot s > a_1^{(2)} \cdot z \geq a_1^{(2)} \cdot D. \quad (4.11)$$

Let

$$C := \{\mathbf{a}_{z+1}, \dots, \mathbf{a}_k\} \setminus \mathbf{A}^* = \{\mathbf{a}_{j_1}, \dots, \mathbf{a}_{j_\mu}\}.$$

Clearly, for every v , $1 \leq v \leq \mu$,

$$a_{j_v}^{(2)} \leq a_1^{(2)} + V(k) \leq a_1^{(2)} + 5D + 5.$$

Since $D > (a_1^{(2)} + 5)/(a_1^{(2)} - 5)$, and by (4.11), we obtain

$$x_s - x_1 \geq a_1^{(2)} \cdot D \geq a_1^{(2)} + 5D + 5 \geq a_{j_v}^{(2)}$$

for every $1 \leq v \leq \mu$.

This implies that

$$\begin{aligned} P_2(\mathbf{A}^{(k)}) &\supset P_2(\mathbf{A}^{(z)}) + P_2(\mathbf{A}^*) + \left\{ \sum_{r=1}^v a_{j_r} \mid 1 \leq v \leq \mu \right\} \\ &\supset \left[T + 1 + a_{j_1}, T + (x_s - x_1) + \sum_{r=1}^{\mu} a_{j_r} \right], \end{aligned} \quad (4.12)$$

i.e. $P_2(\mathbf{A}^{(k)})$ contains an interval of length

$$M := (x_s - x_1) + \sum_{r=2}^{\mu} a_{j_r}^{(2)}.$$

In view of (4.11) and by $a_i^{(2)} \geq a_1^{(2)}$, we have

$$M \geq a_1^{(2)} \cdot z + a_1^{(2)} \cdot (\mu - 1) \geq a_1^{(2)} \cdot \max\{z, \mu - 1\}.$$

Since $k \leq z + 4D + \mu < 5z + \mu$, we obtain $\max\{z, \mu - 1\} > (k - 1)/6 \geq \sqrt{n}/40$. Thus

$$M \geq a_1^{(2)}\sqrt{n}/40. \quad \blacksquare$$

We now turn to the proof of Theorem 4.2.

Define w by

$$H(w) - H(k) = \lfloor H/2 \rfloor.$$

Let us write

$$\mathbf{H}_1 := \{\mathbf{a}_i \mid k < i < w; \mathbf{a}_i \in \mathbf{A}\} \text{ and } \mathbf{H}_2 := \{\mathbf{a}_i \mid w \leq i \leq n; \mathbf{a}_i \in \mathbf{A}\}.$$

By using Lemma 4.3, we obtain that there exists an $U \in \mathbb{N}$ such that for every i , $U < i \leq U + M$, there is a $\mathbf{z}_{j_i} \in P(\mathbf{A}^{(k)})$ for which $z_{j_i}^{(2)} = i$.

Since \mathbf{A} is a simple sequence, it is easy to see that $a_k^{(1)} \leq a_1^{(1)} + k$. This implies that

$$z_{j_i}^{(1)} < a_1^{(1)} \cdot k + k^2 < 2 \cdot k^2 < n/10 \quad (4.13)$$

if n is large enough. Let $\mathbf{z}_{s_1}, \mathbf{z}_{s_2}, \dots, \mathbf{z}_{s_M}$ be a permutation of the sequence $\mathbf{z}_{j_1}, \mathbf{z}_{j_2}, \dots, \mathbf{z}_{j_m}$ such that

$$z_{s_1}^{(1)} \leq z_{s_2}^{(1)} \leq \dots \leq z_{s_M}^{(1)}$$

holds.

By Lemma 4.1 there are $m_1, F \in \mathbb{N}$ such that

$$P(\mathbf{H}_1) \supset \mathcal{H}(F, [H/2]/2, m_1).$$

Let

$$\mathbf{y}_j = (F + 1 + z_{s_M}^{(1)} - z_{s_j}^{(1)}, m_1),$$

for $1 \leq j \leq M$. By (4.13), and since $[H/2]/2 > n/9$, we conclude that $\mathbf{y}_j \in \mathcal{H}(F, [H/2]/2, m_1)$ and

$$\mathbf{z}_{s_j} + \mathbf{y}_j = (F + z_{s_M}^{(1)}, m_1 + s_j)$$

for every $1 \leq j \leq M$.

Thus we obtain

$$\{\mathbf{z}_{s_j} + \mathbf{y}_j \mid j = 1, \dots, M\} \subset \mathcal{V}(F + 1 + z_{s_M}^{(1)}, M, m_1).$$

Furthermore, by using Lemma 4.3 again we obtain that there are $m_2, E \in \mathbb{N}$ such that

$$\mathcal{H}(E, [H/2]/2, m_2) \subset P(\mathbf{H}_2).$$

Since the sets $\mathbf{A}^{(k)}$, \mathbf{H}_1 and \mathbf{H}_2 are pairwise disjoint, applying Lemma 4.2 there are $N, S \in \mathbb{N}$ such that

$$(P(\mathbf{A}^{(k)}) + P(\mathbf{H}_1)) + P(\mathbf{H}_2) \supset \mathbf{R}(N, S, N + M, S + [H/2]/2)$$

holds. This implies that

$$T(n) \geq M \cdot [H/2]/2 \geq (a_1^{(1)} \cdot \sqrt{n}/40) \cdot (n/9) = a_1^{(1)} \cdot n^{\frac{3}{2}}/360 > n^{\frac{3}{2}}/360.$$

Now we are going to construct a set \mathbf{A} for which $T(n) \leq 4n^2$. Let $\mathbf{a}_1 = (0, 0)$ and, for every $n \in \mathbb{N}$, let

$$\mathbf{a}_{n+1} = \mathbf{a}_n + \begin{cases} (1, 0) & \text{if } n \equiv 0 \pmod{2}, \\ (0, 1) & \text{if } n \equiv 1 \pmod{2}. \end{cases}$$

This definition implies that

$$|a_n^{(1)} - a_n^{(2)}| \leq 1.$$

Hence, if $\mathbf{x} \in P(\mathbf{A}^{(n)})$, then

$$|x^{(1)} - x^{(2)}| \leq n. \quad (4.14)$$

Let us assume that, contrary to the assertion, there exists an $\mathbf{R} = \mathbf{R}(a, b, c, d) \subset P(\mathbf{A}^{(n)})$ for which $t(\mathbf{R}) > 4n^2$. This implies that

$$\max\{c - a, d - b\} > 2n$$

and thus $\max\{|b - c|, |b - a|, |d - a|\} > n$. But $\{(a, b), (c, b), (a, d)\} \subset (\mathbf{A}^{(n)})$. This contradicts (4.14) and the proof of the upper bound is completed. \square

PROOF OF THEOREM 4.3. First we are going to prove that there are $\mathbf{x}_1, \mathbf{x}_2, \mathbf{e}_1, \mathbf{e}_2 \in \mathbb{N}^2$ such that $m(\mathbf{e}_i) = m(\mathbf{d}_i)$ and $L(\mathbf{x}_i, \mathbf{e}_i) \subset P(\mathbf{X})$ for $i = 1, 2$.

Let $X_i \subset \mathbb{N}$, $i = 1, 2$, denote the set of integers for which

$$k \in X_i \text{ iff } k \cdot \mathbf{d}_i \in \mathbf{X} \cap L(\mathbf{d}_i). \quad (4.15)$$

This definition implies that

$$m \in P(X_i) \text{ iff } m \cdot \mathbf{d}_i \in P(\mathbf{X} \cap L(\mathbf{d}_i)). \quad (4.16)$$

Furthermore, if $k \cdot \mathbf{d}_i \in L(\mathbf{d}_i) \cap \mathbf{S}_n$, then $k \cdot |\mathbf{d}_i| \leq n$. Hence, by (4.4) and (4.14), we conclude that there are ε' , $0 < \varepsilon' < \varepsilon$, and n_0 such that $X_i(n) > n^{\frac{1}{2} + \varepsilon'}$ for $n > n_0$. Thus, by using Theorem F, we obtain that X_i is subcomplete, i.e. there are $a_{i0}, q_i \in \mathbb{N}$ such that $\{a_{i0} + k \cdot q_i\}_{k=0}^\infty \subset P(X_i)$. Thus, by (4.16), we have

$$\{(a_{i0} + k \cdot q_i) \cdot \mathbf{d}_i\}_{k=0}^\infty \subset P((\mathbf{X} \cap L(\mathbf{d}_i))).$$

Now, if $\mathbf{x}_i = a_{i0} \cdot \mathbf{d}_i$; \mathbf{d}_i ; $\mathbf{e}_i = q_i \cdot \mathbf{d}_i$, then $L(\mathbf{x}_i, \mathbf{e}_i) \subset P(\mathbf{X} \cap L(\mathbf{d}_i))$ for $i = 1, 2$, as we claimed.

Since $m(\mathbf{d}_1) \neq m(\mathbf{d}_2)$, we have $L(\mathbf{x}_1, \mathbf{e}_1) \cap L(\mathbf{x}_2, \mathbf{e}_2) = \emptyset$, and so

$$L := L(\mathbf{x}_1, \mathbf{e}_1) + L(\mathbf{x}_2, \mathbf{e}_2) \subset P(\mathbf{X}).$$

Let $\mathbf{x}_0 = \mathbf{x}_1 + \mathbf{x}_2$. Clearly, $L = \{\mathbf{x}_0 + k \cdot \mathbf{e}_1 + m \cdot \mathbf{e}_2 \mid k, m \in \mathbb{N}\}$. Let $r > |\mathbf{e}_1| + |\mathbf{e}_2|$ and $r = s$. Since the set $\mathbf{A} = \mathbf{X} \setminus (L(\mathbf{d}_1) \cup L(\mathbf{d}_2))$ satisfies (4.3), by Lemmas 4.1 and 4.2 there exist integers a and b such that $\mathbf{R}(a, b, a + r, b + s) \subset P(\mathbf{A})$. Let $\mathbf{w} = (a, b)$ and $\mathbf{y}_0 = \mathbf{x}_0 + \mathbf{w}$. We are going to show that if $\mathbf{z} \in \mathbf{S}(\mathbf{y}_0, \mathbf{e}_1, \mathbf{e}_2)$, then $\mathbf{z} = \mathbf{u} + \mathbf{v}$, where $\mathbf{u} \in L$ and $\mathbf{v} \in P(\mathbf{A})$. Define the integers k and m by

$$m(\mathbf{e}_1) \leq m(\mathbf{z} - (\mathbf{y}_0 + k \cdot \mathbf{e}_1 + m \cdot \mathbf{e}_2)) \leq m(\mathbf{e}_2)$$

and

$$|\mathbf{z} - (\mathbf{y}_0 + k \cdot \mathbf{e}_1 + m \cdot \mathbf{e}_2)| = \min\{|\mathbf{z} - (\mathbf{y}_0 + s \cdot \mathbf{e}_1 + t \cdot \mathbf{e}_2)| : s, t \in \mathbb{N}\}.$$

Since $|\mathbf{z} - (\mathbf{y}_0 + k \cdot \mathbf{e}_1 + m \cdot \mathbf{e}_2)| < |\mathbf{e}_1| + |\mathbf{e}_2| < r$, we obtain that $\mathbf{v} = \mathbf{z} - (\mathbf{y}_0 + k \cdot \mathbf{e}_1 + m \cdot \mathbf{e}_2) \in \mathbf{R}(a, b, a + r, b + r)$. Let $\mathbf{u} = \mathbf{y}_0 + k \cdot \mathbf{e}_1 + m \cdot \mathbf{e}_2$. Thus $\mathbf{u} + \mathbf{v} = \mathbf{z}$, $\mathbf{u} \in L$, $\mathbf{v} \in P(\mathbf{A})$. So we obtain that $\mathbf{S}(\mathbf{y}_0, \mathbf{e}_1, \mathbf{e}_2) \subset P(\mathbf{X})$, i.e. \mathbf{X} is p -complete. \square

REFERENCES

1. T. C. Brown, Is there a sequence of four symbols in which no two adjacent segments are permutations of one another? *Am. Math. Monthly*, **78** (1971), 886–888.
2. J. Folkman, On the representation of integers as sums of distinct terms from a fixed sequence, *Can. J. Math.*, **18** (1966), 643–655.
3. D. A. B. Pleasants, Non-repetitive sequences, *Proc. Camb. Phil. Soc.*, **68** (1970), 267–274.

Received 16 November 1994 and accepted 20 December 1994

NORBERT HEGYVÁRI

Department of Mathematics, Teacher Training School,
L. Eötvös University, Budapest, Hungary
and

Mathematical Institute of the Hungarian Academy of Sciences,
1053 Budapest, Reáltanoda u. 13–15. Hungary